

ITP-93-33E

May 1993

High- Q^2 Elastic ed -scattering and QCD Predictions

A. Kobushkin and A. Syamtomov

*N.N.Bogolyubov Institute for Theoretical Physics,
Ukrainian Academy of Sciences, Kiev 143, Ukraine*

Abstract

In the framework of perturbative QCD it is argued that in the elastic ed -scattering at $Q^2 \sim \text{few } (GeV/c)^2$ the light-cone-frame helicity-flip amplitudes could not be omitted. The obtained $\frac{B}{A}$ ratio of Rosenbluth structure functions is shown to be in a good agreement with experimental data. The high Q^2 behavior of T_{20} is discussed.

In a recent paper, Brodsky and Hiller [1] have made a general analysis of the e.m. properties of spin-one systems proceeding from the gauge theory. They have mentioned that in the analysis of high energy elastic ed -cross-sections at least two momentum scales must be distinguished. The first one is given by the QCD scale $\Lambda_{QCD} \approx 200 MeV/c$ and determines the perturbative QCD regime. In particular, according to arguments of Ref.[2], this scale controls a high Q^2 suppression of the light-cone-frame helicity-flip matrix elements of the deuteron e.m. current

$$\begin{aligned} G_{+0}^+ &\approx a \left(\frac{\Lambda_{QCD}}{Q} \right) G_{00}^+, \\ G_{+-}^+ &\approx b \left(\frac{\Lambda_{QCD}}{Q} \right)^2 G_{00}^+, \end{aligned} \quad (1)$$

where $G_{h'h}^+(Q^2) \equiv \langle p'h' | J^+ | ph \rangle$ are the matrix elements of the e.m. current plus component $J^+ \equiv J^0 + J^3$; $|ph\rangle$ is an eigenstate of momentum p and helicity h ; a and b are some constants.

The second scale is given by the deuteron mass M and has a purely kinematic origin. In Ref.[1] it was argued that in the region

$$Q^2 \gg 2M\Lambda_{QCD} \approx 0.8(GeV/c)^2 \quad (2)$$

helicity-conserving matrix element G_{00}^+ dominates. For the dimensionless ratio $\eta = \frac{Q^2}{4M^2}$ (which appears in expressions for the deuteron e.m. form factors given by (3.7) of Ref.[1]) this condition translates into $\eta \gg \frac{\Lambda_{QCD}}{2M} \approx 0.05$. Omitting the helicity-flip matrix elements G_{+0}^+ and G_{+-}^+ , the following ratios for the charge, magnetic and quadrupole form factors were derived [1]

$$G_C(Q^2) : G_M(Q^2) : G_Q(Q^2) = \left(1 - \frac{2}{3}\eta \right) : 2 : -1. \quad (3)$$

One can note here that the asymptotical relation $G_C \approx \left(\frac{Q^2}{6M^2} \right) G_Q$ of Refs.[2, 3] is reproduced from (3) at considerably high momentum transfer $Q^2 \gg 20(GeV/c)^2$.

In the present paper we would like to mention that the helicity-flip matrix element G_{+0}^+ cannot be neglected. For example, in the light-cone frame defined in Ref.[1] the deuteron magnetic form factor is

$$G_M = \frac{1}{p^+(2\eta + 1)} \left[(2\eta - 1) \frac{G_{+0}^+}{\sqrt{2\eta}} + G_{00}^+ - G_{+-}^+ \right], \quad (4)$$

where p^+ is a plus component of the deuteron momentum. According to (1) the contribution from G_{+-}^+ is suppressed by the factor $\left(\frac{\Lambda_{QCD}}{Q} \right)^2$. But the contribution

from G_{+0}^+ is *enhanced* by the kinematical factor $-\sqrt{\frac{1}{2\eta}}$ at $\eta \ll \frac{1}{2}$ and by the factor $\sqrt{2\eta}$ at $\eta \gg \frac{1}{2}$. It must be also stressed that the signs from G_{+0}^+ contributions are different for the small η -value and large η -value regions. So there may be a point $Q^2 = Q_0^2$ where the contribution from G_{+0}^+ conceals G_{00}^+ matrix element and the magnetic form factor changes its sign. For deuteron this occurs approximately at $2(GeV/c)^2$, what experimental data tell us about [4]. In this case the coefficient a in the expression (1) must be equal to

$$a = \frac{\sqrt{2}Q_0^2 M}{(2M^2 - Q_0^2)\Lambda_{QCD}} \approx 5.08. \quad (5)$$

Adopting this assumption, we obtain the following expressions for the deuteron electromagnetic form factors in the region under consideration

$$\begin{aligned} G_C &= \frac{1}{2p^+(2\eta + 1)} \left(\frac{6M^2 + 5Q_0^2}{6M^2 - 3Q_0^2} - \frac{2}{3}\eta \right) G_{00}^+, \\ G_M &= \frac{1}{p^+(2\eta + 1)} \left(\frac{2\eta - 1}{2\eta} \frac{Q_0^2}{2M^2 - Q_0^2} + 1 \right) G_{00}^+, \\ G_Q &= \frac{1}{2p^+(2\eta + 1)} \left(\frac{1}{\eta} \frac{Q_0^2}{2M^2 - Q_0^2} - 1 \right) G_{00}^+, \end{aligned} \quad (6)$$

and, therefore, the ratio of the Rosenbluth structure functions $\frac{B}{A}$ and tensor analysing power T_{20} become G_{00}^+ -independent. The results of numerical calculations for $\frac{B}{A}$ and T_{20} in comparison with experimental data and the results of Ref.[1] are shown in Figs. 1 and 2, respectively, where Q_0^2 parameter is chosen to be $1.93(GeV/c)^2$.

At very high Q^2 (when $\eta \sim 1$) the ratios (3) are modified to:

$$\begin{aligned} G_C(Q^2) : G_M(Q^2) : G_Q(Q^2) &= \left(\frac{2}{3}\eta - \frac{6M^2 + 5Q_0^2}{6M^2 - 3Q_0^2} \right) : \\ &2 \left(\frac{1 - 2\eta}{2\eta} \frac{Q_0^2}{2M^2 - Q_0^2} - 1 \right) : \left(\frac{1}{\eta} \frac{Q_0^2}{2M^2 - Q_0^2} - 1 \right). \end{aligned} \quad (7)$$

Omitting helicity-flip matrix elements, Brodsky and Hiller [1] showed that at large Q^2 the ratios of deuteron e.m. form factors are identical to those of the point-like spin-one fields. From (7) one sees that such statement remains valid only for $G_C : G_Q$ and only when $\eta \gg 1$.

Some comments should be made about the gross structure of the deuteron e.m. form factors. As was already mentioned, experiment shows that at $Q^2 = Q_0^2$ the magnetic form factor changes its sign. This means that it is negative for $Q^2 > Q_0^2$ and is positive for $Q^2 < Q_0^2$. Therefore the matrix element G_{00}^+ (according to (6)) should

be *negative*. It means (see ratios (7)) that G_C is negative within the interval from $Q^2 \sim 0.8(GeV/c)^2$ (the margin, where the perturbative QCD regime becomes to be open) to $Q^2 \approx 11M^2$. Since G_C is positive at very small Q^2 one can make a conclusion that G_C has to change its sign at $Q \sim$ few hundred MeV/c. Such structure is in a good agreement with conventional meson+nucleon picture predictions (see, e.g. [5]), which are assumed to be valid for $Q^2 \leq 1(GeV/c)^2$. As for the deuteron quadrupole form factor, it is negative in the region where the perturbative QCD is valid, but

$$Q^2 \leq \frac{4M^2Q_0^2}{2M^2 - Q_0^2} \approx 5.32(GeV/c)^2. \quad (8)$$

On the other hand at the origin it is positive. This means that contrary to the meson-nucleon picture, G_Q has a node at some value Q^2 , where perturbative QCD is not yet applicable.

Concluding we should emphasize that our approach is not in contradiction with the G_{00}^+ dominance hypothesis, which, in our opinion, together with available experimental data may allow us to make a reasonable predictions for deuteron electromagnetic form factor behavior at large momentum transfer. We plan to develop an elaborate analysis of one of possible scenarios of such evaluation in a more detailed paper.

The authors are grateful to S. Brodsky for encouragement of this work and to J. Hiller for sending B/A experimental data.

Figure Captions

Figure 1. Comparison of our calculations (solid line) for B/A ratio with experimental data Ref.[4]. The results of calculations of Ref.[1] are plotted by dashed line.

Figure 2. The predictions for T_{20} in our approach (solid line) and the model of Ref.[1] (dashed line) at $\theta = 0^\circ$. Experimental data are from Refs.[6].

References

- [1] S.J. Brodsky and J.R. Hiller, Phys. Rev. D**46** (1992) 2141.
- [2] C.E. Carlson and F. Gross, Phys. Rev. Lett. **53**, (1984) 127.
- [3] C.E. Carlson, Nucl. Phys. A**508** (1990) 481c.
- [4] R.G. Arnold *et al.*, Phys. Rev. Lett. **58**, (1987) 1723; P.E. Bosted *et al.*, Phys. Rev. C**42** (1990) 38; R. Cramer *et al.*, Z. Phys. C**29** (1985) 513; S. Platchkov *et al.*, Nucl. Phys. A**510**, (1990) 740.
- [5] M. Gari and H. Hyuga, Nucl. Phys. A**264** (1976) 409.
- [6] I. The *et al.*, Phys. Rev. Lett. **67**, (1991) 173; R. Gilman *et al.*, *ibid.***65**, (1990) 1733; B.B. Voitsekhovshii *et al.*, Pis'ma Zh. Eksp. Teor. Fiz. **43**, (1986) 567 [JETP Lett. **43**, (1986) 733]; M.E. Schulze *et al.*, Phys. Rev. Lett. **52**, (1984) 597.




Fig. 1




Fig. 2